
Thesis on Dysentery. Edwin P. Aller of Philade In admitted March 10. 1821

amita March 10. 1821



## On Dysentery

Sheer is implanted in the human breast; a desire to impact something which is new and interesting. In the fursivit of our ancient, and highly humanable profession, research affects ample opportunity for the gradification of this laudable ambiliant. The writer had intended to have taken up and stablished, in some measure, the character of a newly-intereduced active of the Materiae Medical; but his views have been interrupted, by the insuperable difficulties have a student has be knowned; thought do how his he hands he trusts he will seer the insuperable difficulties have a student has be knowned; the unity asks at your hands he trusts he will seer the inergies of his limited understanding, to become a worthy obile of the School which reased him, and of the Sountry which gover him booth f.

The disease which he has therefore chosen as the subject of his dissertation is, the Dysonting. the most

In treating of the above malady, a student labours under many disadvantages, for independent left under his care, he is, (undappily for the advancement of under his care, to is, (undappily for the advancement of under his care, centineally perplaced with the conflicting destrines of the wistest and most judicious of the profession. When the wistest and most judicious of the profession. When the wistest and most judicious of the profession when his therefore, he enters upon the duties of, his station, he finds himself, necessitated to depend, almost solely, upon his come judgment. Their soperiories is placed before him. So is with him to determine, whether he will declare him to lite the left, or, whether, he will declare him tell independent, and act accordingly.

At present, I shall not attempt to pass judgment on the actual merits or demerits of the several authors whom I shall call in to my wide, but shall be satisfied withe making a compilation from the practice of those whom I consider as the most inlightened and shillful; not free suming to comment much when them.

Typentery is a disease, which consists in a fever, accompanied with frequent mucous, and sometimes, bloody stools; attended with flatulence, termina, tenesmus, and a retention of the natural faces.

Whis disease may occur at any season of the year, thought generally, it commences its march in the United States about the latter and of the seventh menth (Suly) or the beginning of the 8th. me: August, and continues until the Swellth menth December.

Similed, as is my experience in this disease, sham neverthely dishored to doub! the validity of the assution that a specific contagion, is to be considered, as always the remote cause there.

The fact of its depending upon a morbid state of the Atmosphere.

for its dissemination, Scensider as clearly preven by the high est authorities which I have consulted.

e 4the many instances are recorded wherein Depending has attacked in succession several members of the same family it does not calidate the conclusion, that it is communicated by specific infection from one to another, and for the following reasons.

First. That there is a peculiarity in the atmosphere precise pring

<sup>&</sup>quot; bullens First Sines far: 1074.

poing the body to disease, and determining what that aisease may be, by look setuation. He no no fine Toyunteres he vailing in high grounds, while sowers obtain in the callies in their immediate vicinite.

Meendly. What the whencene soft greater in Cambes and Societals, does not melitate against their who down its inher tens nature, recourse we find it appearing after tellines march is coherent to incliment weather, and at that sensen if the year when there are sudden changes from heat to, cold, all trees are bredisposine course.

But, it man be asked, who was it attach, one and unother, and unother in an apparently general succession, until the whole an under jet influence? Who wish that when patients are early removed from these large assemblies the disease is more mild, and that when the healthy are separated from the rest, it disappears? "Breause, a predisposition existings committee at a superfict in the rest, it disappears? "Breause, a predisposition existings committee at the rest with what at first and sinchere, renders it still more so, and need-sorily increases its virulence: and vice versa.



the much them, in consideration of the injections character, of Bysently. In subject of my healtern Tooler to Before Chapman, Brilester, Galdwell; Mydenham, Williet to to

Mystenham, and his followers, have denominated this disense a fobris introversa; a four sturned in upon the bowels; but 3' um inclined to adopt the opinion of many highly respectable breatitions of the present day, that primarily it is a gastrie, but, that ultimately it is a gastrie, but, that ultimately it breams can intestinal disease.

Courring at the same line of the year with automoral intermetants and remittents, it is with them frequently com-

It is remented a concrete in it the relieving or emeritary on the so the reason and a service to concrete an electronic of the reason and a service of the mouth and combined with named and remitting. These complimes are sometimed encounted in a short time, to movid arterial actions descreted in the head secretic things of the below the secretic things of the bead secretic things of the bead secretic things of the bead secretic things of the things and the things and the secretic beautiful the continuing, were converted berooklibs, becomes now full and strong and

White his between delivered in the Bush's dy down ham? The University of Bush of Paris of Pharmas Role dec 1. oak 111.

\* Bush Bush St. Some Higher Winer Systember Set.



exerciating bains harrass the treets.

Mu other instances, the luin in the securis comes on unaccompanies by any apparent febrile affection, and is succeeded by a line smust, and frequent, but small discharges, from the intestines, consisting chiefly of an intermistence of blood and muses, Instahr with these samplems, the appliete for second becomes an uncertain but nausen and comiting rarely cover. The an uncertain but hoften the flattent is uttacked with fover, in a more or less violent degree, and withit the lemina and tenesmus brewne more severe, and the steels more frequent.

The matter evacuated by stool is very varied in its ah tearance: semilimes it is entirely mucous, exhibiting that form of disease, by some called moons mucous; and by other, assuring alvair at other times the discharge is solely sanguineous; but generally, we find an intermiature of blood and mucous.

"The sivels are all along distinguished by a peculiar small differing from that of commen exact months; the always offension. It wants the six bet towards the



end, when the intestines begin to mortify, the fator, is cadaverous and intelevable.

Pouthers have neliced membrane-like substances, supplied to be abrasiens of the villens cout of the intestines. Also, while sebaccous masses are sometimes discharged per anum resembling such and have thereford been called corpora funquia. The nature of these substances, however, had I believe, not get been satisfactorily determined.

We have stated that the transations to steel are very frauent, yet this exerctions are seldom natural, and appear in small industrial masses termed bearing, which being evided, affects temperary relief to the agencyed policies, from termina stemmus. Some speak of worms being discharged during the course of the disease to so, wamme, but this may be considered as an adventitions circumstance. I procedultie and and shippars sim of veries are semetimes attendants!

Then the bowels become affected with gangrene, the palient ne linear complains of bain or tensmus, the hulse sinks; deterium and come generally supervine : in some instances, newerer, the mind remains saw and correfled until death.



Wissections present the intestines, harticularly the octen and return) in a highly inflamed and semetimes mertilied; state; also present, scirchosdies, alcoration, contractions, and numerous. The perituaum and other severings of the abdomainments instances, have likewess an inflammatery appearance.

Ther much interesting matter as regards dest morten we searches Arefer to Pringle & Clipherns.

From what has been stated me may with sasely conclude, that the Agrentery chiefly provails at the same season of the year with Intermettent and Remittent Greens, and that it a rises from the same causes, in general, by which they are from the same

## Causes.

· Weach has been said respecting the causes of Dysentery), but nothing satisfactory has been offend, by then practitioners, whose



opinions of have read; there is good reason, however, to estimit, that among the free disposing causes are to be enumerated, is trems in the weathers from het and day, to evel and damp or rainy; debiting of body, either in continuous of free rains discussed, the under observation of the mind, or interviewed the reading or disching, and excessive fategues. Whe excitent causes are, the application of construct herstradium and hossibly, acoustives a suppression of the acoustimed herstradium and bossibly, acoustive irritating, substances taken into the stemach.

## - · !realment! -

Whatever the cause of Dysentery may be, the indications of our uppear to be the same, namely;

1st. The verceme the inflammation stricture of the intestines, is in other words. We obtain intent Jain and fiver.

I not. it breserve the towels soluble until all shasm, pain , and fiver have disappeared .

But. He restere the healthy action of the skin.

11h. To restore the line of the ulimentary conul.



These indications are best unswered by the following Memedies.

1st. Bleedletting.

The exterience of the most eminent and successful Startitioned of Course pe, and of our can country in fractions in Afficients. Sufficiently warrants the recurrence to this excellent antiphlodistic and unidown: Indicate upon over this incide of reduced there, it would seem to have the respectively of their she quantitie of the course is to be requisited in more on the deserve; from and obstinces of the stricters, than on the halfs, which in all diseases of an influential nature of feeling the thinks via is an uncertain in draw of the state of the rotant of the latter the stricted from and extending, some to the country of the 20 encountry of the state of the state of the started of the s

A melies, which must be safely new accommissioned. Of their class of remedies I would be sire to Intermedian Swaterizatum, telieving that no specific Intellegenteric vistue is associable to Sheenewankar And also on account of the grouter Grapherette, and Mathartic, as well as Courte effect of Varter Emeric.

Bia. Laxatives. I say laxatives, for in the irritable state



of the affected harte it is presumable that fungatives esticially of the deaths kind, would be likely to renew the train efficients since sumptions. Among the mild and officient aloine concesses, the Chrem Stevini is deserving of primary attention, and should be viven in the dose of half an ensure; in a little Someonade, on While hall, is very 2nd hour puril two or three facual concentions now been precured. If the Oil should not be retained on the ottemach, an excellent substitute is, a solution of Mathhas Soda, in a strong infusion of Mathhas Loda, in a strong infusion of Mamemile, sav 2 concess of the state, in a string infusion: a wine deasyle to a given at intervals of, 2 hours while faces be freely evacuated.

Another valuable medicine is the Brotechleride (Hebmariate) of Reveny, in combination with Satch, to which the Chiem Anisi may be added by way of Carminative, thus,

R. bal. ppl. Zfs Puli Julap. Vij Cl. Anisipec.

Nitee, et ctivie: in bht is Mumat ager emam emni bikeris uomee bene eferaverint. This herover beniters in the fugative ) Ms auxiliaries, the fatient may cirink freely of barley-water; "This valuable muciture, or that of Jum Irabic or Flax seed Few, with the addition of Laudanum; forms an Enema. Seculially calculated to bloviate the extreme irritability, and consequent Uncomes so characteristic of the cisease.

Metted butter fore from sall regrancially in the briter tien of 5 will or half a hint, as an injection is highly sal utare!

linsudrea, thin calin gruel, balm tea, of mucilage of Mippery.

The any of the above laxatives a landamen may be added, accoming to the judgment of the physician, invase of source pain of the breeks, without fear of charities until the pendian sugarative was a line freely used landier until the pendian sugarative vacuations have yielded to what I might call landed stocks; it will be proper to divert the ottermination to the skin, by Diraphretics, the best of which Scensides the Bilovis Specaculanha bomposities or Divers Powders in deux of fix vacury 4th hours continued for one or 2 day, individual third feet by this hort facts by the thick the fitted by the survey 4th hours of surv

5 h is not to be forgotton that a large Epispustic over the abstemen is one of the indispensables, in cases where great irritability prevails. If the stemach cannot be painfies by the usual internal premailes, a blister must be applied over the epigastrie region.

Pek. Having thus, as will generally be the case, overcome the disease, it remains, to restore the tene of the alimentary canal.

This is effected by the class of Firsts, which ought to be used with caution, commencing with the milder and changing to there of a more powerful kind. I would first use the Perpentaria and bort: Aurant: , must the Rhubarb in small dails, then bort: Cinchona. During the use of these, the Nutrientia perperty come in, such as animal broths, jellies, cysters, seft boiled eigs, feels-, with Porter, Me, Madeira, seeund Cyder, and the like.

In aid of the tonics, riding on horse back, when practi-

cable, ought not to be omitted.

Thomas Page waited March 17, 1021